




coffee&health

from the institute for scientific information on coffee



**Fact or Fiction**



Coffee is one of the most widely researched food items in the world. Overall, the large and growing body of scientific research shows that coffee, when drunk in moderation i.e. 3-5 regular cups a day, is perfectly safe for the majority of healthy adults.

However, there are still many misconceptions about coffee and health which can lead to confusion about whether coffee consumption can be enjoyed as part of a healthy, balanced diet. You may come across these in your daily practice as your patients or colleagues ask you for advice.

Based on the latest scientific research, this document, which has been developed for health care and other professional audiences, serves as a quick-reference guide to respond to some of the most frequently asked questions about coffee and health.

**Drinking coffee is bad for health**

**Drinking coffee does not increase the risk of cardiovascular disease**

**Pregnant women should stop drinking coffee**

**Drinking coffee helps improve sports performance**

**Coffee is dehydrating**

**Drinking coffee keeps me alert and helps me concentrate**

**Coffee can become addictive**

**Drinking coffee in the afternoon or evening does not always disrupt sleep**

**Decaffeinated coffee is healthier than regular coffee**

## **Drinking coffee is bad for health**

### **FICTION**

- Regular coffee drinking can be part of a healthy, balanced diet and lifestyle, and moderate coffee consumption i.e. 3-5 regular cups a day, is safe for most individuals with no adverse effects
- Scientific evidence also suggests that moderate coffee consumption may actually offer a number of benefits. For example, the European Food Safety Authority recently stated that caffeine improves both sports performance (endurance exercise) and increases attention and alertness <sup>(1,2)</sup>
- However some individuals may choose to switch to decaffeinated coffee e.g. pregnant women, those sensitive to the stimulant effects of caffeine late afternoon/early evening. For these individuals, decaffeinated coffee provides an alternative so they can still enjoy the taste and aroma of coffee



## Drinking coffee does not increase the risk of cardiovascular disease

### FACT

- Moderate coffee consumption is not linked to an increased risk of cardiovascular problems such as heart disease, heart attacks, irregular heart beat or high blood pressure <sup>(3,4)</sup>
- Research also suggests that, in both men and women, drinking coffee in moderation may reduce their risk of stroke, but no firm conclusion has yet been drawn <sup>(5)</sup>
- Coffee's effect on cholesterol levels is largely dependent on the method of brewing. Filtered coffee is not associated with a significant increase in cholesterol levels, while boiled coffee can raise cholesterol levels <sup>(6)</sup>

## Pregnant women should stop drinking coffee

### FICTION

- Studies from the last decade clearly report that moderate caffeine consumption, including that from coffee, is not a matter of concern for a healthy pregnancy <sup>(7)</sup>. Pregnant women can still enjoy a couple of cups of regular coffee a day
- Health authorities recommend that pregnant women should limit their daily caffeine intake to 200-300mg from all sources <sup>(8)</sup>. A regular cup of caffeinated coffee contains approximately 80-100mg of caffeine
- Overall, well-conducted scientific studies in humans have shown no adverse effects on the fetus if a pregnant woman consumes a moderate amount of caffeine from coffee or other caffeinated beverages

## Drinking coffee helps improve sports performance

### FACT

- The effects of coffee consumption on sports performance are linked to the caffeine in coffee, rather than to coffee itself
- The European Food Safety Authority (EFSA) recently stated that a cause and effect relationship has been established for caffeine intake and increased endurance performance, endurance capacity, and a reduction in perceived exertion <sup>(1)</sup>. Caffeine is effective at doses of 3-4mg/kg
- Caffeine may moderate central fatigue and influence ratings of perceived exertion, pain and levels of vigour, all of which are likely to lead to improvements in performance <sup>(9)</sup>



## Coffee is dehydrating

### FICTION

- While there is some indication of a mild, short-term diuretic effect of caffeine, this effect is not strong enough to counter-balance the benefits of fluid intake from coffee drinking
- Scientific evidence looking at the effects of caffeine on fluid balance does not support a significant diuretic effect of caffeine <sup>(11)</sup>
- Coffee drinking in moderation contributes to our daily fluid intake and does not lead to dehydration, or significant loss of body fluid <sup>(12)</sup>

## Drinking coffee keeps me alert and helps me concentrate

### FACT

- The caffeine in coffee is well known for its stimulating effects, which have scientifically proven benefits on mental performance
- The European Food Safety Authority (EFSA) recently stated that a 75mg serving of caffeine (the amount found in approximately one regular cup of coffee) increases both attention and alertness <sup>(2)</sup>
- The stimulant effects of a regular cup of coffee are observed between 15 – 45 minutes after consumption and normally last for about four hours <sup>(10)</sup>

## Coffee can become addictive

### FICTION

- While caffeine in coffee is a mild central nervous system stimulant, recent scientific studies using brain scans suggest that moderate coffee drinkers do not develop a physical dependence to caffeine <sup>(13,14,15)</sup>
- Some studies suggest that removing caffeine from the diet suddenly may lead to mild, temporary withdrawal symptoms, like headache, in some individuals. These symptoms can be avoided by a gradual reduction of caffeine intake from the diet over time <sup>(16)</sup>
- It is likely that people continue to drink coffee because they enjoy its taste and aroma, and recognise it as a behavioural stimulant; and not because of any addictive qualities of caffeine



## Drinking coffee in the afternoon or evening does not always disrupt sleep

### FACT

- Some people who are sensitive to caffeine find that the mild stimulation of coffee, consumed late in the afternoon or shortly before going to bed, may delay the time it takes them to fall asleep and/or affects their sleep overall. Other people consume caffeinated drinks during the evening and have no problems falling asleep
- Anyone who is sensitive to the stimulant effects of caffeine can enjoy decaffeinated coffee during the afternoon and evening instead <sup>(17)</sup>
- It should be noted that there are many other factors, including noise, temperature, and discomfort, that may affect how long it takes someone to get to sleep <sup>(18)</sup>.

## Decaffeinated coffee is healthier than regular coffee

### FICTION

- There are some circumstances when individuals may be advised to, or choose to, switch to decaffeinated coffee. Pregnant women, for example, are advised to limit their caffeine intake to 200mg - 300mg per day <sup>(8)</sup> and also those very sensitive to caffeine. For those individuals, decaffeinated coffee provides an alternative so they can still enjoy the taste and aroma of coffee
- In addition, some people find that the mild stimulant effect of caffeine consumed late in the afternoon or shortly before going to bed affects their sleep. In their case, it is well advised to switch to decaffeinated coffee during the afternoon and evening <sup>(17)</sup>
- In other cases, drinking caffeinated coffee has actually been shown to have some health benefits. For example, the European Food Safety Authority recently stated that caffeine improves both sports performance (endurance exercise) and increases attention and alertness <sup>(1,2)</sup>

For more information, please visit [www.coffeeandhealth.org](http://www.coffeeandhealth.org)



## References

1. EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA) (2011). Scientific Opinion on the substantiation of health claims related to caffeine and increase in physical performance during short-term high-intensity exercise (ID 737, 1486, 1489), increase in endurance performance (ID 737, 1486), increase in endurance capacity (ID 1488) and reduction in the rated perceived exertion/effort during exercise (ID 1488, 1490) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. EFSA Journal;9(4):2053. Available at [www.efsa.europa.eu/en/efsajournal/pub/2053.htm](http://www.efsa.europa.eu/en/efsajournal/pub/2053.htm)
2. EFSA (2011). Scientific Opinion on the substantiation of health claims related to caffeine and increased alertness (ID 736, 1101, 1187, 1485, 1491, 2063, 2103), and increased attention (ID 736, 1485, 1491, 2375) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. EFSA Journal;9(4):2054. Available at: [www.efsa.europa.eu/en/efsajournal/pub/2054.htm](http://www.efsa.europa.eu/en/efsajournal/pub/2054.htm)
3. Lopez-Garcia E et al. (2011) Coffee consumption and mortality in women with cardiovascular disease. *Am J Clin Nutr*; 94 (4):1113-1126.
4. Klatsky AL et al (2011). Coffee, caffeine, and risk of hospitalization for arrhythmias. *The Permanente Journal*; 15(3): 1519-25.
5. S C Larrson SC & Orsini N (2011). Coffee consumption and risk of stroke: A dose-response meta-analysis of prospective studies, *Am J Epidemiol*; doi: 10.1093/aje/kwr226
6. Jee SH et al (2001). Coffee consumption and serum lipids: a meta-analysis of randomized controlled trials. *Am j Epidemiol*; 153:353-62.
7. Peck JD et al (2010). A review of the epidemiological evidence concerning the reproductive health effects of caffeine consumption: a 2000-2009 update. *Food Chem Toxicol*; 48:2549-76.
8. American College of Obstetrician and Gynaecologists (2010). Moderate caffeine consumption during pregnancy. Committee opinion no. 462. *Obstet Gynecol*; 116:467-8.
9. Davis JK et al (2009). Caffeine and anaerobic performance: ergogenic value and mechanisms of action. *Sports Med*; 39:813-32.
10. Fredholm BB et al (1999). Actions of caffeine in the brain with special reference to factors that contribute to widespread use. *Pharmacol Rev*; 51:83-133.
11. Popkin BM et al (2006). A new proposed guidance system for beverage consumption in the United States. *Am J Clin Nutr*, 83, 529-542.
12. Kolasa KM et al (2009). Hydration and health promotion. *Nutrition Today* 44, 190-203.
13. Nehlig A et al (2000). Dose-response study of caffeine effects on cerebral functional activity with a specific focus on dependence. *Brain Res*; 858:71-77.
14. Acquas E et al (2002). Differential effects of caffeine on dopamine and acetylcholine transmission in brain areas of drug-naive and caffeine-pretreated rats. *Neuropsychopharmacology*; 27:182-193.
15. De Luca MA et al (2007). Caffeine and accumbens shell dopamine. *J Neurochem*; 103:157-163.
16. Nehlig A (2004). Are we dependent on coffee and caffeine: an update. In Nehlig A, ed. *Coffee, Tea, Chocolate and the Brain*. Boca Raton, FL: CRC Press; 133-146.
17. Sin CWM, et al (2008). Systematic review on the effectiveness of caffeine abstinence on the quality of sleep. *J Clin Nursing*; 18:13-21.
18. Carrier J et al (2009). Effects of caffeine on daytime recovery sleep: A double challenge to the sleep-wake cycle in aging. *Sleep Med*;10:1016-24.